Actavis

NDC 0591-**2981**-35

Compare to the active ingredient in Abreva®**

Docosanol Cream, 10%

Cold Sore/Fever Blister Treatment

Non-prescription Medicine to Shorten Healing Time*

SAMPLE

Cold Sore Treatment For Topical Use Only

> Net Wt. 2a (.07 oz)



Distributed by: Actavis Pharma, Inc. Parsippany, NJ 07054 USA 29810117 R1118 DP22662 AZU ant m absiM



with, manufactured by, or produced by the makers or owners of Abreva®. This product is not affiliated

duration of symptoms. рир әшіз биірәү sore medicine to shorten *Non-prescription cold

NO COATING

Monday-Friday. **Questions?** 1-888-838-2872 between 9 am and 5 pm ET,

benzyl alcohol, light mineral oil, propylene glycol, purfiled water, sucrose distearate, sucrose stearate Inactive ingredients

- do not freeze
- Other information

 store at 20° to 25°C (68° to 77°F)
- children under 12 years: ask a doctor
- . up in gently but completely . use 5 times a day until healed sore/fever blister (tingle)

 • early treatment ensures the best results
- wash hands before and after applying cream
 wash hands before and after applying cream
 - adults and children 12 years or over: Directions

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. Drug Facts (continued)

sore is not hea**l**ed within 10 days Stop use and ask a doctor if • your cold sore gets worse or the cold

When using this product • apply only to the affected areas • do not use in or near the eyes • avoid applying directly inside your mouth o do not share this product with anyone. This may spread the infection. Do not use • if you are allergic to any ingredient in this product

For external use only

If an allergic reaction occurs, stop use and seek medical help right away. SHOCK • RASH
 Include: • NIVES This product may cause a severe allergic reaction. Symptoms may noclude: • hives • facial swelling • wheezing/difficulty breathing

Milergy Alert Warnings

 tingling, pain, burning, and/or itcning shortens healing time and duration of symptoms:

USGS • treats cold sores/fever blisters on the face or lips

Cold sore/fever blister treatment Docosanol 10% Active ingredient Purpose

Drug Facts

Assisting in serifor tull product uses, directions and warnings,

Docosanol Cream, 10%

SAMPLE

Thank you for purchasing Docosanol Cream. We hope that since Docosanol Cream shortens the healing time of cold sores, it helps you look and feel better faster. Please follow insert directions for proper use and refer to the information below.

You are not alone

An estimated one-in-five Americans is affected by cold sores each year.

Most people who experience them get two to three cold sore outbreaks each year.

Just before and during an outbreak, the cold sore infection can be easily passed on to others. It is especially important to be aware of early warning signs that a cold sore is coming, and to be sure to avoid close physical contact when you have one.

Getting to the Root of the Problem

A blister, or cluster of blisters around your mouth, is caused by the cold sore virus. After the cold sore has healed, the virus stays dormant in your body but can be reactivated by everyday triggers like stress, sunlight, extreme hot or cold weather, colds and flu, lack of sleep, or menstruation.

The Sooner, the Better

A typical cold sore outbreak can last as long as seven to ten days. The sooner you begin using Docosanol Cream, the more effective the medicine will be.

So for best results, use Docosanol Cream at the very start of the cold sore – when you feel the first sign of the tingle, redness, bump or itch.

Docosanol Cream is different from other over-the-counter cold sore products. Only Docosanol Cream contains 10% docosanol, which helps to shorten the healing time of cold sores and the duration of symptoms including tingling, pain, itching and burning.

Frequently Asked Questions:

How is Docosanol Cream different from the over-the-counter cold sore remedies I have already been using?

Docosanol Cream is a non-prescription cold sore medicine that can actually shorten healing time and the duration of symptoms. The sooner you start using Docosanol Cream, the sooner you can begin healing your cold sore.

Who can use Docosanol Cream?

Adults and children 12 years and over can use Docosanol Cream.

How often can I apply Docosanol Cream?

Docosanol Cream should be used five times a day until the cold sore is healed. Apply it directly to the cold sore at the first sign of a tingle, redness, bump or itch for best results.

Can I apply cosmetics on top of Docosanol Cream?

Yes, cosmetics such as lipstick may be applied over Docosanol Cream. However, use a separate applicator, like a cotton swab, to apply cosmetics over an unhealed cold sore to avoid spreading the infection. For best results, remove any cosmetics prior to applying Docosanol Cream.

Made in the USA
Distributed by: Actavis Pharma, Inc.
Parsippany, NJ 07054 USA
Revised – March 2016



Docosanol Cream, 10%

For Topical Use Only
Net Wt. 2g (.07 oz)

Retain insert for full product uses, directions and warnings.

Made in the USA

SAMPLE

Distributed by: Actavis Pharma, Inc. Parsippany, NJ 07054 USA 29810117 R0718 236834-02



