

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use **EMTRICITABINE and TENOVIFOR DISOPROXIL FUMARATE TABLETS** safely and effectively. See full prescribing information for **EMTRICITABINE and TENOVIFOR DISOPROXIL FUMARATE TABLETS**. **EMTRICITABINE and TENOVIFOR DISOPROXIL FUMARATE TABLETS**, for oral use Initial U.S. Approval: 2004

WARNING: POSTTREATMENT ACUTE EXACERBATION OF HEPATITIS B AND RISK OF DRUG RESISTANCE WITH USE OF EMTRICITABINE AND TENOVIFOR DISOPROXIL FUMARATE TABLETS FOR HIV-1 PRE-EXPOSURE PROPHYLAXIS (PrEP) IN UNDIAGNOSED EARLY HIV-1 INFECTION

See full prescribing information for complete boxed warning.

- Severe acute exacerbations of hepatitis B (HBV) have been reported in HIV-infected individuals who have discontinued emtricitabine and tenofovir disoproxil fumarate tablets. Hepatic function should be monitored closely in these individuals who discontinue emtricitabine and tenofovir disoproxil fumarate tablets. If appropriate, anti-hepatitis B therapy may be warranted. (5.1)
- Emtricitabine and tenofovir disoproxil fumarate tablets used for HIV-1 PrEP should be prescribed to individuals confirmed to be HIV-negative immediately prior to initiating and at least every 3 months during use. Drug-resistant HIV-1 variants have been identified with use of emtricitabine and tenofovir disoproxil fumarate tablets for HIV-1 PrEP following undetected acute HIV-1 infection. Do not initiate emtricitabine and tenofovir disoproxil fumarate tablets for HIV-1 PrEP if signs or symptoms of acute HIV infection are present unless negative infection status is confirmed. (5.2)

RECENT MAJOR CHANGES

Indications and Usage
HIV-1 Pre-Exposure Prophylaxis (PrEP) (1.2) 06/20/20
Dosage and Administration
HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP (2.2) 06/20/20
Warnings and Precautions
Severe Acute Exacerbation of Hepatitis B and Risk of Sexually Transmitted Infections, Including HIV-1, and Development of HIV-1 Resistance When Emtricitabine and Tenofovir Disoproxil Fumarate Tablets are Used for HIV-1 PrEP (5.2) 06/20/20
Immune Reconstitution Syndrome (5.4) 06/20/20

INDICATIONS AND USAGE
HIV-1 Treatment (1.1)
Emtricitabine and tenofovir disoproxil fumarate tablets are a two-drug combination of emtricitabine (FTC) and tenofovir disoproxil fumarate (TDF), both HIV-1 nucleoside analog reverse transcriptase inhibitors, and is indicated for:
in combination with other antiretroviral agents for the treatment of HIV-1 infection in adults and pediatric patients weighing at least 17 kg.

HIV-1 PrEP (1.2)
Emtricitabine and tenofovir disoproxil fumarate tablets are indicated in at-risk adults and adolescents weighing at least 35 kg for pre-exposure prophylaxis (PrEP) to reduce the risk of sexually acquired HIV-1 infection. Individuals must have a negative HIV-1 test immediately prior to initiating emtricitabine and tenofovir disoproxil fumarate tablets for HIV-1 PrEP.

DRUG INTERACTIONS
Tenofovir disoproxil fumarate increases didanosine concentrations. Dose reduction and close monitoring of didanosine toxicity are warranted.
Coadministration decreases atazanavir concentrations. When coadministered with emtricitabine and tenofovir disoproxil fumarate tablets, use atazanavir with ritonavir. (7.2)

WARNINGS AND PRECAUTIONS
Testing: Prior to or when initiating emtricitabine and tenofovir disoproxil fumarate tablets test for hepatitis B virus infection. Prior to initiation and during use of emtricitabine and tenofovir disoproxil fumarate tablets, on a clinically appropriate schedule, assess serum creatinine, estimated creatinine clearance, uric acid, and urine protein in all individuals. In individuals with chronic kidney disease, also assess serum phosphorus. (2.4)

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

133 mg/200 mg, or 167 mg/250 mg based on body weight) once daily taken orally with or without food. (2.4)
2.4 Recommended dosage in renally impaired HIV-1 infected adult patients:
o Creatinine clearance (CrCl) 30–49 mL/min: 1 tablet every 48 hours. (2.6)
o CrCl below 30 mL/min or hemodialysis: emtricitabine and tenofovir disoproxil fumarate tablets are not recommended. (2.2)

HIV-1 Pre-Exposure Prophylaxis (PrEP)
Recommended dosage in HIV-1 uninfected adults and adolescents weighing at least 35 kg: One emtricitabine and tenofovir disoproxil fumarate tablet (containing 200 mg of FTC and 300 mg of TDF) once daily taken orally with or without food. (2.5)
2.2 Recommended dosage in renally impaired HIV-1 uninfected individuals: emtricitabine and tenofovir disoproxil fumarate tablets are not recommended in HIV-1 uninfected individuals if CrCl is below 30 mL/min. (2.6)

DOSAGE FORMS AND STRENGTHS
Tablets: 200 mg/300 mg of emtricitabine and tenofovir disoproxil fumarate, respectively. (3)
CONTRAINDICATIONS
Emtricitabine and tenofovir disoproxil fumarate tablets for HIV-1 PrEP are contraindicated in individuals with unknown or positive HIV-1 status. (4)

WARNINGS AND PRECAUTIONS
Severe Acute Exacerbation of Hepatitis B and Risk of Sexually Transmitted Infections, Including HIV-1, and Development of HIV-1 Resistance When Emtricitabine and Tenofovir Disoproxil Fumarate Tablets are Used for HIV-1 PrEP
Lactic Acidosis/Severe Hepatomegaly with Steatosis
Discontinue emtricitabine and tenofovir disoproxil fumarate tablets in individuals who develop symptoms or laboratory findings suggestive of lactic acidosis or pronounced hepatomegaly. (5.6)

ADVERSE REACTIONS
In HIV-1 uninfected patients, the most common adverse reactions (incidence greater than or equal to 10%) are diarrhea, nausea, fatigue, headache, dizziness, depression, insomnia, abnormal dreams, and rash. (6.1)
In HIV-1 uninfected adults in PrEP trials, adverse reactions that were reported in more than 2% of emtricitabine and tenofovir disoproxil fumarate tablet participants and more frequently than by placebo recipients were headache, abdominal pain, and weight decreased. (6.1)

SUSPECTED ADVERSE REACTIONS, CONTACT YOUR HEALTHCARE PROVIDER OR VISIT www.fda.gov/medwatch

DRUG INTERACTIONS
Tenofovir disoproxil fumarate increases didanosine concentrations. Dose reduction and close monitoring of didanosine toxicity are warranted.
Coadministration decreases atazanavir concentrations. When coadministered with emtricitabine and tenofovir disoproxil fumarate tablets, use atazanavir with ritonavir. (7.2)

WARNINGS AND PRECAUTIONS
Testing: Prior to or when initiating emtricitabine and tenofovir disoproxil fumarate tablets test for hepatitis B virus infection. Prior to initiation and during use of emtricitabine and tenofovir disoproxil fumarate tablets, on a clinically appropriate schedule, assess serum creatinine, estimated creatinine clearance, uric acid, and urine protein in all individuals. In individuals with chronic kidney disease, also assess serum phosphorus. (2.4)

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

DOSAGE AND ADMINISTRATION

2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection or for HIV-1 PrEP
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

2.3 Recommended Dosage for Treatment of HIV-1 Infection in Adults and Pediatric Patients Weighing at Least 35 kg
2.4 Recommended Dosage for Treatment of HIV-1 Infection in Pediatric Patients Weighing at Least 17 kg and Able to Swallow a Tablet

2.5 Recommended Dosage for Treatment of HIV-1 Infection in Pediatric Patients Weighing 17 kg to Less Than 35 kg
2.6 Dosage Adjustment in Individuals with Renal Impairment

2.7 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.8 Dosage Adjustment in Individuals with Renal Impairment

2.9 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.10 Dosage Adjustment in Individuals with Renal Impairment

2.11 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.12 Dosage Adjustment in Individuals with Renal Impairment

2.13 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.14 Dosage Adjustment in Individuals with Renal Impairment

2.15 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.16 Dosage Adjustment in Individuals with Renal Impairment

2.17 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.18 Dosage Adjustment in Individuals with Renal Impairment

2.19 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.20 Dosage Adjustment in Individuals with Renal Impairment

2.21 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.22 Dosage Adjustment in Individuals with Renal Impairment

2.23 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.24 Dosage Adjustment in Individuals with Renal Impairment

2.25 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.26 Dosage Adjustment in Individuals with Renal Impairment

2.27 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.28 Dosage Adjustment in Individuals with Renal Impairment

2.29 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.30 Dosage Adjustment in Individuals with Renal Impairment

2.31 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.32 Dosage Adjustment in Individuals with Renal Impairment

2.33 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.34 Dosage Adjustment in Individuals with Renal Impairment

2.35 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.36 Dosage Adjustment in Individuals with Renal Impairment

2.37 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.38 Dosage Adjustment in Individuals with Renal Impairment

2.39 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.40 Dosage Adjustment in Individuals with Renal Impairment

2.41 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.42 Dosage Adjustment in Individuals with Renal Impairment

2.43 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.44 Dosage Adjustment in Individuals with Renal Impairment

2.45 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.46 Dosage Adjustment in Individuals with Renal Impairment

2.47 Recommended Dosage for HIV-1 PrEP in Adults and Adolescents Weighing at Least 35 kg
2.48 Dosage Adjustment in Individuals with Renal Impairment

Autimmune disorders (such as Graves' disease, polymyositis, Guillain-Barré syndrome, and autoimmune hepatitis) have also been reported to occur in the setting of immune reconstitution; however, the time to onset is more variable and can also occur many months after initiation of treatment.
Bone Loss and Mineralization Defects
In clinical trials in HIV-1 uninfected adults and in a clinical trial of HIV-1 uninfected individuals, TDF (a component of emtricitabine and tenofovir disoproxil fumarate tablets) was associated with slightly greater decreases in bone mineral density (BMD) and increases in biochemical markers of bone metabolism, suggesting increased bone turnover relative to comparators. (See *Adverse Reactions* (6.1)). Serum parathyroid hormone levels and 1,25 Vitamin D levels were also higher in subjects receiving TDF.

Concomitant use of emtricitabine and tenofovir disoproxil fumarate tablets and close monitoring of renal function are recommended in all patients with estimated creatinine clearance 30 to 49 mL/min. (See *Adverse Reactions* (6.1)).
In HIV-1 uninfected individuals, decreases in BMD were observed in the TDF-treated HIV-1 uninfected pediatric subjects as compared to the control groups. Similar trends were observed in adolescents with chronic kidney disease, also assessed serum phosphorus. (See *Adverse Reactions* (6.1)).
In all pediatric trials, skeletal growth (height) appeared to be unaffected.
The effects of TDF-associated changes in BMD and biochemical markers on long-term bone health and future fracture risk are unknown. Assessment of BMD should be considered for adult and pediatric patients who have a history of fractures and/or osteoporosis and are taking TDF. However, although the effect of supplementation with calcium and vitamin D was not studied, such supplementation may be beneficial. If bone abnormalities are suspected, appropriate consultation should be obtained.

Mineralization Defects
Lactic Acidosis/Severe Hepatomegaly with Steatosis
Discontinue emtricitabine and tenofovir disoproxil fumarate tablets in individuals who develop symptoms or laboratory findings suggestive of lactic acidosis or pronounced hepatomegaly. (5.6)

ADVERSE REACTIONS
In HIV-1 uninfected patients, the most common adverse reactions (incidence greater than or equal to 10%) are diarrhea, nausea, fatigue, headache, dizziness, depression, insomnia, abnormal dreams, and rash. (6.1)
In HIV-1 uninfected adults in PrEP trials, adverse reactions that were reported in more than 2% of emtricitabine and tenofovir disoproxil fumarate tablet participants and more frequently than by placebo recipients were headache, abdominal pain, and weight decreased. (6.1)

SUSPECTED ADVERSE REACTIONS, CONTACT YOUR HEALTHCARE PROVIDER OR VISIT www.fda.gov/medwatch

DRUG INTERACTIONS
Tenofovir disoproxil fumarate increases didanosine concentrations. Dose reduction and close monitoring of didanosine toxicity are warranted.
Coadministration decreases atazanavir concentrations. When coadministered with emtricitabine and tenofovir disoproxil fumarate tablets, use atazanavir with ritonavir. (7.2)

WARNINGS AND PRECAUTIONS
Testing: Prior to or when initiating emtricitabine and tenofovir disoproxil fumarate tablets test for hepatitis B virus infection. Prior to initiation and during use of emtricitabine and tenofovir disoproxil fumarate tablets, on a clinically appropriate schedule, assess serum creatinine, estimated creatinine clearance, uric acid, and urine protein in all individuals. In individuals with chronic kidney disease, also assess serum phosphorus. (2.4)

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

INDICATIONS AND USAGE
1.1 Treatment of HIV-1 Infection
1.2 HIV-1 Pre-Exposure Prophylaxis (PrEP)
2 DOSAGE AND ADMINISTRATION
2.1 Testing Prior to Initiation of Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for Treatment of HIV-1 Infection
2.2 HIV-1 Screening for Individuals Receiving Emtricitabine and Tenofovir Disoproxil Fumarate Tablets for HIV-1 PrEP

Autimmune disorders (such as Graves' disease, polymyositis, Guillain-Barré syndrome, and autoimmune hepatitis) have also been reported to occur in the setting of immune reconstitution; however, the time to onset is more variable and can also occur many months after initiation of treatment.
Bone Loss and Mineralization Defects
In clinical trials in HIV-1 uninfected adults and in a clinical trial of HIV-1 uninfected individuals, TDF (a component of emtricitabine and tenofovir disoproxil fumarate tablets) was associated with slightly greater decreases in bone mineral density (BMD

How do I tell my healthcare provider before taking emtricitabine and tenofovir disoproxil fumarate tablets?

Before taking emtricitabine and tenofovir disoproxil fumarate tablets, tell your healthcare provider about all of your medical conditions, including if you:

- have liver problems, including HBV infection
- have kidney problems or receive kidney dialysis treatment
- have bone problems
- are pregnant or plan to become pregnant. It is not known if emtricitabine and tenofovir disoproxil fumarate can harm your unborn baby. Tell your healthcare provider if you become pregnant during treatment with emtricitabine and tenofovir disoproxil fumarate tablets.

Pregnancy Registry: There is a pregnancy registry for people who take emtricitabine and tenofovir disoproxil fumarate tablets during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. Talk with your healthcare provider about how you can take part in this registry.

- are breastfeeding or plan to breastfeed. Emtricitabine and tenofovir disoproxil fumarate can pass to your baby in your breast milk.
- Do not breastfeed if you have HIV-1 or if you think you have recently become infected with HIV-1 because of the risk of passing HIV-1 to your baby.
- If you take emtricitabine and tenofovir disoproxil fumarate tablets for HIV-1 PrEP, talk with your healthcare provider about the best way to feed your baby.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Some medicines may interact with emtricitabine and tenofovir disoproxil fumarate tablets. Keep a list of your medicines and show it to your healthcare provider and pharmacist when you get a new medicine.

You can ask your healthcare provider or pharmacist for a list of medicines that interact with emtricitabine and tenofovir disoproxil fumarate tablets.

Do not start a new medicine without telling your healthcare provider. Your healthcare provider can tell you if it is safe to take emtricitabine and tenofovir disoproxil fumarate tablets with other medicines.

How should I take emtricitabine and tenofovir disoproxil fumarate tablets?

- Take emtricitabine and tenofovir disoproxil fumarate tablets exactly as your healthcare provider tells you to take it.
- If you take emtricitabine and tenofovir disoproxil fumarate tablets to treat HIV-1 infection, you need to take other HIV-1 medicines. Your healthcare provider will tell you what medicines to take and how to take them.
- Take emtricitabine and tenofovir disoproxil fumarate tablets 1 time each day with or without food.
- Children who take emtricitabine and tenofovir disoproxil fumarate tablets are prescribed a lower strength tablet than adults. Children should swallow the emtricitabine and tenofovir disoproxil fumarate tablet. Tell your healthcare provider if your child cannot swallow the tablet, because they may need a different HIV-1 medicine.
- Your healthcare provider will change the dose of emtricitabine and tenofovir disoproxil fumarate tablets as needed based on your child's weight.

Do not change your dose or stop taking emtricitabine and tenofovir disoproxil fumarate tablets without first talking with your healthcare provider. Stay under a healthcare provider's care when taking emtricitabine and tenofovir disoproxil fumarate tablets.

Do not miss a dose of emtricitabine and tenofovir disoproxil fumarate tablets.

If you take too many emtricitabine and tenofovir disoproxil fumarate tablets, call your healthcare provider or go to the nearest hospital emergency room right away.

When your emtricitabine and tenofovir disoproxil fumarate tablets supply starts to run low, get more from your healthcare provider or pharmacy.

- If you are taking emtricitabine and tenofovir disoproxil fumarate tablets for treatment of HIV-1, the amount of virus in your blood may increase if the medicine is stopped for even a short time. The virus will develop resistance to emtricitabine and tenofovir disoproxil fumarate tablets and become harder to treat.
- If you take too many emtricitabine and tenofovir disoproxil fumarate tablets, call your healthcare provider or go to the nearest hospital emergency room right away.
- When your emtricitabine and tenofovir disoproxil fumarate tablets supply starts to run low, get more from your healthcare provider or pharmacy.

Caution: If you are taking emtricitabine and tenofovir disoproxil fumarate tablets for treatment of HIV-1, the amount of virus in your blood may increase if the medicine is stopped for even a short time. The virus will develop resistance to emtricitabine and tenofovir disoproxil fumarate tablets and become harder to treat.

Caution: If you take too many emtricitabine and tenofovir disoproxil fumarate tablets, call your healthcare provider or go to the nearest hospital emergency room right away.

Caution: When your emtricitabine and tenofovir disoproxil fumarate tablets supply starts to run low, get more from your healthcare provider or pharmacy.

Caution: Do not change your dose or stop taking emtricitabine and tenofovir disoproxil fumarate tablets without first talking with your healthcare provider. Stay under a healthcare provider's care when taking emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: Do not miss a dose of emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: If you take too many emtricitabine and tenofovir disoproxil fumarate tablets, call your healthcare provider or go to the nearest hospital emergency room right away.

Caution: When your emtricitabine and tenofovir disoproxil fumarate tablets supply starts to run low, get more from your healthcare provider or pharmacy.

Caution: Do not change your dose or stop taking emtricitabine and tenofovir disoproxil fumarate tablets without first talking with your healthcare provider. Stay under a healthcare provider's care when taking emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: Do not miss a dose of emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: If you take too many emtricitabine and tenofovir disoproxil fumarate tablets, call your healthcare provider or go to the nearest hospital emergency room right away.

Caution: When your emtricitabine and tenofovir disoproxil fumarate tablets supply starts to run low, get more from your healthcare provider or pharmacy.

Caution: Do not change your dose or stop taking emtricitabine and tenofovir disoproxil fumarate tablets without first talking with your healthcare provider. Stay under a healthcare provider's care when taking emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: Do not miss a dose of emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: If you take too many emtricitabine and tenofovir disoproxil fumarate tablets, call your healthcare provider or go to the nearest hospital emergency room right away.

Caution: When your emtricitabine and tenofovir disoproxil fumarate tablets supply starts to run low, get more from your healthcare provider or pharmacy.

Caution: Do not change your dose or stop taking emtricitabine and tenofovir disoproxil fumarate tablets without first talking with your healthcare provider. Stay under a healthcare provider's care when taking emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: Do not miss a dose of emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: If you take too many emtricitabine and tenofovir disoproxil fumarate tablets, call your healthcare provider or go to the nearest hospital emergency room right away.

Caution: When your emtricitabine and tenofovir disoproxil fumarate tablets supply starts to run low, get more from your healthcare provider or pharmacy.

Caution: Do not change your dose or stop taking emtricitabine and tenofovir disoproxil fumarate tablets without first talking with your healthcare provider. Stay under a healthcare provider's care when taking emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: Do not miss a dose of emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: If you take too many emtricitabine and tenofovir disoproxil fumarate tablets, call your healthcare provider or go to the nearest hospital emergency room right away.

Caution: When your emtricitabine and tenofovir disoproxil fumarate tablets supply starts to run low, get more from your healthcare provider or pharmacy.

Caution: Do not change your dose or stop taking emtricitabine and tenofovir disoproxil fumarate tablets without first talking with your healthcare provider. Stay under a healthcare provider's care when taking emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: Do not miss a dose of emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: If you take too many emtricitabine and tenofovir disoproxil fumarate tablets, call your healthcare provider or go to the nearest hospital emergency room right away.

Caution: When your emtricitabine and tenofovir disoproxil fumarate tablets supply starts to run low, get more from your healthcare provider or pharmacy.

Caution: Do not change your dose or stop taking emtricitabine and tenofovir disoproxil fumarate tablets without first talking with your healthcare provider. Stay under a healthcare provider's care when taking emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: Do not miss a dose of emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: If you take too many emtricitabine and tenofovir disoproxil fumarate tablets, call your healthcare provider or go to the nearest hospital emergency room right away.

Caution: When your emtricitabine and tenofovir disoproxil fumarate tablets supply starts to run low, get more from your healthcare provider or pharmacy.

Caution: Do not change your dose or stop taking emtricitabine and tenofovir disoproxil fumarate tablets without first talking with your healthcare provider. Stay under a healthcare provider's care when taking emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: Do not miss a dose of emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: If you take too many emtricitabine and tenofovir disoproxil fumarate tablets, call your healthcare provider or go to the nearest hospital emergency room right away.

Caution: When your emtricitabine and tenofovir disoproxil fumarate tablets supply starts to run low, get more from your healthcare provider or pharmacy.

Caution: Do not change your dose or stop taking emtricitabine and tenofovir disoproxil fumarate tablets without first talking with your healthcare provider. Stay under a healthcare provider's care when taking emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: Do not miss a dose of emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: If you take too many emtricitabine and tenofovir disoproxil fumarate tablets, call your healthcare provider or go to the nearest hospital emergency room right away.

Caution: When your emtricitabine and tenofovir disoproxil fumarate tablets supply starts to run low, get more from your healthcare provider or pharmacy.

Caution: Do not change your dose or stop taking emtricitabine and tenofovir disoproxil fumarate tablets without first talking with your healthcare provider. Stay under a healthcare provider's care when taking emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: Do not miss a dose of emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: If you take too many emtricitabine and tenofovir disoproxil fumarate tablets, call your healthcare provider or go to the nearest hospital emergency room right away.

Caution: When your emtricitabine and tenofovir disoproxil fumarate tablets supply starts to run low, get more from your healthcare provider or pharmacy.


Caution: Do not change your dose or stop taking emtricitabine and tenofovir disoproxil fumarate tablets without first talking with your healthcare provider. Stay under a healthcare provider's care when taking emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: Do not miss a dose of emtricitabine and tenofovir disoproxil fumarate tablets.

Caution: If you take too many emtricitabine and tenofovir disoproxil fumarate tablets, call your healthcare provider or go to the nearest hospital emergency room right away.


Caution: When your emtricitabine and tenofovir disoproxil fumarate tablets supply starts to run low, get more from your healthcare provider or pharmacy.

It has the following structure:



FTC is a white to off-white crystalline powder with a solubility of approximately 112 mg/mL in water at 25°C. The partition coefficient (log P) for emtricitabine is -0.43 and the pKa is 2.65.

Tenofovir Disoproxil Fumarate: TDF is a fumarate acid salt of the bis-isopropoxycarbonyloxymethyl ester derivative of tenofovir. The chemical name of tenofovir DF is 9-(R)-[2-[(bis[(isopropoxy)carbonyloxy]-methoxy]phosphoryl)methyl]propane]diolamine fumarate. It has the following structural formula:



Tenofovir disoproxil fumarate is a white to off-white crystalline powder with a solubility of 13.4 mg/mL in water at 25°C. The partition coefficient (log P) for tenofovir disoproxil is 1.23 and the pKa is 3.73. All dosages are expressed in terms of TDF except where otherwise noted.

Emtricitabine and tenofovir disoproxil fumarate tablets are for oral administration. Each film-coated tablet contains 200 mg of emtricitabine and 300 mg of tenofovir DF (which is equivalent to 245 mg of tenofovir disoproxil), as active ingredients. The tablets also include the following inactive ingredients: croscarmellose, FD&C Blue #2 aluminum lake, lactose monohydrate, magnesium stearate, mannitol, polyethylene glycol, polyvinyl alcohol, povidone, talc, and titanium dioxide.

12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
Emtricitabine and tenofovir disoproxil fumarate tablets are a fixed-dose combination of antiviral drugs FTC and TDF (see Microbiology (12.2)).

12.3 Pharmacokinetics
Emtricitabine and Tenofovir Disoproxil Fumarate Tablets: One emtricitabine and tenofovir disoproxil fumarate tablet was comparable to one FTC capsule (200 mg) plus one TDF tablet (300 mg) following single-dose administration to fasting healthy subjects (N = 38).

Emtricitabine: The pharmacokinetic properties of FTC are summarized in Table 8. Following oral administration of FTC, FTC is rapidly absorbed with peak plasma concentrations occurring at 1 to 2 hours postdose. Less than 4% of FTC binds to human plasma proteins in vitro, and the binding is independent of concentration over the range of 0.02 to 200 mcg/mL. Following administration of radiolabeled FTC, approximately 86% is recovered in the urine and 13% is recovered as metabolites. The metabolites of FTC include 3'-sulfoxide diastereomers and their glucuronic acid conjugates.

Emtricitabine is eliminated by a combination of glomerular filtration and active tubular secretion. Following a single oral dose of FTC, the plasma FTC half-life is approximately 10 hours.

Tenofovir Disoproxil Fumarate: The pharmacokinetic properties of TDF are summarized in Table 8. Following oral administration of TDF, mean tenofovir serum concentrations are achieved within 1 to 2 hours. Less than 0.7% of tenofovir binds to human plasma proteins in vitro, and the binding is independent of concentration over the range of 0.01 to 25 mcg/mL. Approximately 70 to 80% of the intravenous dose of tenofovir is recovered as unchanged drug in the urine. Tenofovir is eliminated by a combination of glomerular filtration and active tubular secretion. Following a single oral dose of TDF, the terminal elimination half-life of tenofovir is approximately 17 hours.

Table 8 Single Dose Pharmacokinetic Parameters for FTC and Tenofovir in Adults^a

	FTC	Tenofovir
Fasted Oral Bioavailability ^b (%)	92 (83.1 to 106.4)	25 (N/C to 45.0)
Plasma Terminal Elimination Half-Life ^c (hr)	10 (7.4 to 15.0)	17 (12.0 to 25.7)
C _{max} ^d (mcg/mL)	1.8 ± 0.72 ^e	0.30 ± 0.09
AUC ^f (mcg·hr/mL)	10.0 ± 3.12 ^e	2.29 ± 0.69
CL/F ^g (mL/min)	302 ± 94	1043 ± 315
CL _{renal} ^h (mL/min)	213 ± 89	243 ± 33

a. N/C = Not calculated.
b. Median (range).
c. Mean ± SD.
d. Data presented as steady state values.

Effects of Food on Oral Absorption
Emtricitabine and tenofovir disoproxil fumarate tablets may be administered with or without food. Administration of emtricitabine and tenofovir disoproxil fumarate tablets following a high fat meal (784 kcal; 49 grams of fat) or a light meal (373 kcal; 8 grams of fat) delayed the time of tenofovir C_{max} by approximately 0.75 hour. The mean increases in tenofovir AUC and C_{max} were approximately 55% and 15%, respectively, when administered with a high fat or light meal, compared to administration in the fasted state. In previous safety and efficacy trials, TDF (tenofovir) was taken under fast conditions. FTC systemic exposures (AUC and C_{max}) were unaffected when emtricitabine and tenofovir disoproxil fumarate tablets were administered with either a high fat or a light meal.

Specific Populations
Paediatrics: No pharmacokinetic differences due to race have been identified following the administration of FTC.

Tenofovir Disoproxil Fumarate: There were insufficient numbers from racial and ethnic groups other than Caucasian to adequately determine potential pharmacokinetic differences among these populations following the administration of TDF.

Gender: Emtricitabine and Tenofovir Disoproxil Fumarate: FTC and tenofovir pharmacokinetics are similar in male and female subjects.

Pediatric Patients
Treatment of HIV-1 Infection: The pharmacokinetic data for tenofovir and FTC following administration of emtricitabine and tenofovir disoproxil fumarate tablets in pediatric subjects weighing 17 kg and above are not available. The dosage recommendations of emtricitabine and tenofovir disoproxil fumarate tablets in this population are based on the dosage recommendations of FTC and TDF in this population. Refer to the EMTRIVA and VIREAD prescribing information for pharmacokinetic information on the individual products in pediatric patients.

Geriatric Patients
Pharmacokinetics of FTC and tenofovir have not been fully evaluated in the elderly (65 years of age and older).

Patients with Renal Impairment
The pharmacokinetics of FTC and tenofovir are altered in subjects with renal impairment (see Warnings and Precautions (5.2)). In adult subjects with creatinine clearance below 50 mL/min, AUC and AUC₀₋₁₂ of FTC and tenofovir were increased. No data are available to make dosage recommendations in pediatric patients with renal impairment.

Patients with Hepatic Impairment
The pharmacokinetics of tenofovir following a 300 mg dose of TDF have been studied in non-HIV-infected subjects with moderate to severe hepatic impairment. There were no substantial alterations in tenofovir pharmacokinetics in subjects with hepatic impairment compared with unpaired subjects. The pharmacokinetics of emtricitabine and tenofovir disoproxil fumarate tablets or FTC have not been studied in subjects with hepatic impairment; however, FTC is not significantly metabolized by liver enzymes, so the impact of liver impairment should be limited.

Assessment of Drug Interactions
The steady state pharmacokinetics of FTC and tenofovir were unaffected when FTC and TDF were administered together versus each agent doses alone.

In vitro studies and clinical pharmacokinetic drug-drug interaction trials have shown that the potential for CYP mediated interactions involving FTC and tenofovir with other medicinal products is low.

TDF is a substrate of P-glycoprotein (P-gp) and breast cancer resistance protein (BCRP) transporters. When TDF is coadministered with an inhibitor of these transporters, an increase in absorption may be observed.

No clinically significant drug interactions have been observed between FTC and famciclovir, indinavir, stavudine, TDF and zidovudine (Tables 9 and 10). Similarly, no clinically significant drug interactions have been observed between TDF and efavirenz, methadone, nelfinavir, oral contraceptives, rifabutin, or zalcitabine in trials conducted in healthy volunteers (Tables 11 and 12).

Table 9 Drug Interactions: Changes in Pharmacokinetic Parameters for FTC in the Presence of the Coadministered Drug^a

Coadministered Drug	Dose of Coadministered Drug (mg)	FTC Dose (mg)	N	% Change of FTC Pharmacokinetic Parameters ^b (90% CI)		
				C _{max}	AUC	C _{min}
TDF	300 once daily x 7 days	200 once daily x 7 days	17	↔	↔	↔
Zidovudine	300 twice daily x 7 days	200 once daily x 7 days	27	↔	↔	↔
Indinavir	800 x 1 x 7 days	200 x 1 x 12	12	↔	↔	NA
Famciclovir	500 x 1 x 7 days	200 x 1 x 12	12	↔	↔	NA
Stavudine	40 x 1 x 7 days	200 x 1 x 12	6	↔	↔	NA

a. All interaction trials conducted in healthy volunteers.
b. ↑ = Increase; ↔ = No Effect; NA = Not Applicable

Table 10 Drug Interactions: Changes in Pharmacokinetic Parameters for Coadministered Drug in the Presence of FTC^a

Coadministered Drug	Dose of Coadministered Drug (mg)	FTC Dose (mg)	N	% Change of Coadministered Drug Pharmacokinetic Parameters ^b (90% CI)		
				C _{max}	AUC	C _{min}
TDF	300 once daily x 7 days	200 once daily x 7 days	17	↔	↔	↔
Zidovudine	300 twice daily x 7 days	200 once daily x 7 days	27	↑ 17	↑ 13	↔
Indinavir	800 x 1 x 7 days	200 x 1 x 12	12	↔	↔	NA
Famciclovir	500 x 1 x 7 days	200 x 1 x 12	12	↔	↔	NA
Stavudine	40 x 1 x 7 days	200 x 1 x 12	6	↔	↔	NA

a. All interaction trials conducted in healthy volunteers.
b. ↑ = Increase; ↔ = No Effect; NA = Not Applicable

Table 11 Drug Interactions: Changes in Pharmacokinetic Parameters for Tenofovir in the Presence of the Coadministered Drug

Coadministered Drug	Dose of Coadministered Drug (mg)	N	% Change of Tenofovir Pharmacokinetic Parameters ^b (90% CI)		
			C _{max}	AUC	C _{min}
Atazanavir ^c	400 once daily x 14 days	33	1.4 (1.8 to 2.0)	1.24 (1.21 to 1.28)	7.22 (1.15 to 3.0)
Atazanavir/ Ritonavir ^c	300/100 once daily	12	34 (20 to 51)	37 (30 to 45)	29 (21 to 36)
Darunavir/ Ritonavir ^d	300/100 twice daily	12	1.24 (0.8 to 1.42)	1.22 (1.10 to 1.35)	1.37 (1.07 to 1.57)
Indinavir	800 three times daily x 7 days	13	1.4 (1.3 to 1.33)	1.35	↔
Ledipasvir/ Sofosbuvir ^{e,1}	90/400 once daily x 10 days	24	1.47 (1.37 to 1.58)	1.35 (1.29 to 1.42)	1.47 (1.38 to 1.57)
Ledipasvir/ Sofosbuvir ^{e,2}	90/400 once daily x 14 days	23	1.64 (1.54 to 1.74)	1.59 (1.42 to 1.70)	1.59 (1.49 to 1.70)
Ledipasvir/ Sofosbuvir ^{e,3}	90/400 once daily x 14 days	15	1.73 (1.56 to 1.94)	1.78 (1.77 to 1.82)	1.63 (1.32 to 1.97)
Ledipasvir/ Sofosbuvir ^{e,4}	90/400 once daily x 14 days	14	1.32 (1.20 to 1.39)	1.31 (1.10 to 1.50)	1.91 (1.74 to 2.10)
Ledipasvir/ Sofosbuvir ^{e,5}	90/400 once daily x 10 days	29	1.61 (1.51 to 1.72)	1.65 (1.59 to 1.71)	1.115 (1.05 to 1.128)
Lopinavir/ Ritonavir	400/100 twice daily x 14 days	24	1.46 (1.32 to 1.61)	1.52 (1.37 to 1.67)	1.51 (1.37 to 1.66)
Saquinavir/ Ritonavir	1000/100 twice daily x 14 days	35	↔	↔	↔
Sofosbuvir ^e	400 single dose	16	1.25 (1.8 to 1.45)	↔	↔
Sofosbuvir/ Velpatasvir ^e	400/100 once daily	24	1.44 (1.33 to 1.55)	1.40 (1.34 to 1.46)	1.84 (1.76 to 1.92)
Sofosbuvir/ Velpatasvir ^e	400/100 once daily	30	1.46 (1.39 to 1.54)	1.40 (1.34 to 1.46)	1.84 (1.61 to 1.79)
Sofosbuvir/ Velpatasvir/ Voxilaprevir ^e	400/100/100 x 29	19	1.36 (1.61 to 1.61)	1.38 (1.42 to 1.45)	1.38 (1.56)
Tacrolimus	0.05 mg/kg twice daily x 7 days	21	1.13 (1.10 to 1.27)	1.2	1.7
Tipranavir/ Ritonavir ⁶	500/100 twice daily	22	1.32 (1.13 to 1.3)	1.2 (1.2 to 1.17)	1.2
	750/200 twice daily (23 doses)	20	1.38 (1.46 to 1.29)	1.2 (1.6 to 1.10)	1.14 (1.1 to 1.27)

a. Subjects received VIREAD 300 mg once daily.
b. Increase = ↑, Decrease = ↓, No Effect = ↔.
c. Reyataz Prescribing Information.
d. Prezista Prescribing Information.
e. Emtricitabine and Tenofovir Disoproxil Fumarate Tablets: Staggered administration (12 weeks apart) provided similar results.
f. Comparison based on exposures when administered as atazanavir/ritonavir + FTC/TDF.
g. Comparison based on exposures when administered as darunavir/ritonavir + FTC/TDF.
h. Comparison based on exposures when administered as atazanavir/ritonavir + FTC/TDF.
i. Study conducted with ATRIPLA (efavirenz/FTC/TDF) coadministered with HARVONI.
j. Study conducted with COMPLERA (FTC/rilpivirine/TDF) coadministered with HARVONI.
k. Study conducted with emtricitabine and tenofovir disoproxil fumarate tablets (FTC/TDF) + dolutegravir coadministered with HARVONI.
l. Study conducted with ATRIPLA coadministered with SOVALDI® (sofosbuvir).
m. Study conducted with COMPLERA coadministered with EPLUSUA, coadministration with EPLUSUA also results in comparable increases in tenofovir exposures when TDF is administered as ATRIPLA, STRIBILD, emtricitabine and tenofovir disoproxil fumarate tablets + atazanavir/ritonavir, or emtricitabine and tenofovir disoproxil fumarate tablets + darunavir/ritonavir.
n. Administered as atazanavir + FTC/TDF.
o. Comparison based on exposures when administered as darunavir + ritonavir + FTC/TDF.
p. Study conducted with additional voxilaprevir 100 mg to achieve voxilaprevir exposures expected in HCV-infected patients coadministered with HARVONI.
q. Aptivus Prescribing Information.

No effect on the pharmacokinetic parameters of the following coadministered drugs was observed with emtricitabine and tenofovir disoproxil fumarate tablets: abacavir, didanosine (buffered tablets), FTC, efavirenz, and lamivudine.

Table 12 Drug Interactions: Changes in Pharmacokinetic Parameters for Coadministered Drug in the Presence of Tenofovir

Coadministered Drug	Dose of Coadministered Drug (mg)	N	% Change of Coadministered Drug Pharmacokinetic Parameters ^b (90% CI)		
			C _{max}	AUC	C _{min}